LECTURE PRESENTATIONS For CAMPBELL BIOLOGY, NINTH EDITION

Jane B. Reece, Lisa A. Urry, Michael L. Cain, Steven A. Wasserman, Peter V. Minorsky, Robert B. Jackson

Chapter 10

Photosynthesis

Lectures by Erin Barley Kathleen Fitzpatrick

Overview: The Process That Feeds the Biosphere

- Photosynthesis is the process that converts solar energy into chemical energy
- Directly or indirectly, photosynthesis nourishes almost the entire living world

- Autotrophs sustain themselves without eating anything derived from other organisms
- Autotrophs are the producers of the biosphere, producing organic molecules from CO₂ and other inorganic molecules
- Almost all plants are photoautotrophs, using the energy of sunlight to make organic molecules


Figure 10.1



- Photosynthesis occurs in plants, algae, certain other protists, and some prokaryotes
- These organisms feed not only themselves but also most of the living world



Figure 10.2



- Heterotrophs obtain their organic material from other organisms
- Heterotrophs are the consumers of the biosphere
- Almost all heterotrophs, including humans, depend on photoautotrophs for food and O₂

- The Earth's supply of fossil fuels was formed from the remains of organisms that died hundreds of millions of years ago
- In a sense, fossil fuels represent stores of solar energy from the distant past




Concept 10.1: Photosynthesis converts light energy to the chemical energy of food


- Chloroplasts are structurally similar to and likely evolved from photosynthetic bacteria
- The structural organization of these cells allows for the chemical reactions of photosynthesis

Chloroplasts: The Sites of Photosynthesis in Plants

- Leaves are the major locations of photosynthesis
- Their green color is from chlorophyll, the green pigment within chloroplasts
- Chloroplasts are found mainly in cells of the mesophyll, the interior tissue of the leaf
- Each mesophyll cell contains 30–40 chloroplasts

- CO₂ enters and O₂ exits the leaf through microscopic pores called stomata
- The chlorophyll is in the membranes of thylakoids (connected sacs in the chloroplast); thylakoids may be stacked in columns called grana
- Chloroplasts also contain stroma, a dense interior fluid







Tracking Atoms Through Photosynthesis: *Scientific Inquiry*


 Photosynthesis is a complex series of reactions that can be summarized as the following equation:

 $6 \text{ CO}_2 + 12 \text{ H}_2\text{O} + \text{Light energy} \rightarrow \text{C}_6\text{H}_{12}\text{O}_6 + 6 \text{ O}_2 + 6 \text{ H}_2\text{O}$

Resp. $C_{GH_{12}} = 760_2 - 760_2 + 6H_2 = 760_2 + 76$ tEnergy


The Splitting of Water

 Chloroplasts split H₂O into hydrogen and oxygen, incorporating the electrons of hydrogen into sugar molecules and releasing oxygen as a byproduct



Photosynthesis as a Redox Process

- Photosynthesis reverses the direction of electron flow compared to respiration
- Photosynthesis is a redox process in which H₂O is oxidized and CO₂ is reduced
- Photosynthesis is an endergonic process; the energy boost is provided by light





The Two Stages of Photosynthesis: A Preview


- Photosynthesis consists of the light reactions (the *photo* part) and Calvin cycle (the *synthesis* part)
- The light reactions (in the thylakoids)
 - Split H20 take e
 - Release O₂
 - Reduce **NADP**⁺ to NADPH
 - Generate ATP from ADP by? Electron transport photophosphorylation Sand ATP synthese





- The Calvin cycle (in the stroma) forms sugar cyce from CO₂, using ATP and NADPH
- The Calvin cycle begins with carbon fixation, incorporating CO₂ into organic molecules











Concept 10.2: The light reactions convert solar energy to the chemical energy of ATP and NADPH


- Chloroplasts are solar-powered chemical factories
- Their thylakoids transform light energy into the chemical energy of ATP and NADPH

The Nature of Sunlight

- Light is a form of electromagnetic energy, also called electromagnetic radiation
- Like other electromagnetic energy, light travels in rhythmic waves
- Wavelength is the distance between crests of waves
- Wavelength determines the type of electromagnetic energy

- The **electromagnetic spectrum** is the entire range of electromagnetic energy, or radiation
- Visible light consists of wavelengths (including those that drive photosynthesis) that produce colors we can see
- Light also behaves as though it consists of discrete particles, called photons


Figure 10.7



Photosynthetic Pigments: The Light Receptors

- Pigments are substances that absorb visible light
- Different pigments absorb different wavelengths
- Wavelengths that are not absorbed are reflected or transmitted
- Leaves appear green because chlorophyll reflects and transmits green light






- A spectrophotometer measures a pigment's ability to absorb various wavelengths
- This machine sends light through pigments and measures the fraction of light transmitted at each wavelength

Figure 10.9


TECHNIQUE



- An absorption spectrum is a graph plotting a pigment's light absorption versus wavelength
- The absorption spectrum of chlorophyll a suggests that violet-blue and red light work best for photosynthesis
- An action spectrum profiles the relative effectiveness of different wavelengths of radiation in driving a process




RESULTS



- The action spectrum of photosynthesis was first demonstrated in 1883 by Theodor W. Engelmann
- In his experiment, he exposed different segments of a filamentous alga to different wavelengths
- Areas receiving wavelengths favorable to photosynthesis produced excess O₂
- He used the growth of aerobic bacteria clustered along the alga as a measure of O₂ production


- Chlorophyll *a* is the main photosynthetic pigment
- Accessory pigments, such as chlorophyll b, broaden the spectrum used for photosynthesis
- Accessory pigments called carotenoids absorb excessive light that would damage chlorophyll

Figure 10.11



Excitation of Chlorophyll by Light

- When a pigment absorbs light, it goes from a ground state to an excited state, which is unstable
- When excited electrons fall back to the ground state, photons are given off, an afterglow called fluorescence
- If illuminated, an isolated solution of chlorophyll will fluoresce, giving off light and heat






(a) Excitation of isolated chlorophyll molecule

(b) Fluorescence

© 3011 Pearson Education, Inc.


A Photosystem: A Reaction-Center Complex Associated with Light-Harvesting Complexes

- A photosystem consists of a reaction-center complex (a type of protein complex) surrounded by light-harvesting complexes
- The light-harvesting complexes (pigment molecules bound to proteins) transfer the energy of photons to the reaction center




(a) How a photosystem harvests light

@ 2011 Peason Education, Inc.




(b) Structure of photosystem II



(a) How a photosystem harvests light

© 2011 Peanson Education, Inc.

Thylakoid membrane



(b) Structure of photosystem II

@ 2011 Pearson Education, Inc.

- A primary electron acceptor in the reaction center accepts excited electrons and is reduced as a result
- Solar-powered transfer of an electron from a chlorophyll a molecule to the primary electron acceptor is the first step of the light reactions


- There are two types of photosystems in the thylakoid membrane
- Photosystem II (PS II) functions first (the numbers reflect order of discovery) and is best at absorbing a wavelength of 680 nm
- The reaction-center chlorophyll a of PS II is called P680

- Photosystem I (PS I) is best at absorbing a wavelength of 700 nm
- The reaction-center chlorophyll a of PS I is called P700


Linear Electron Flow

- During the light reactions, there are two possible routes for electron flow: cyclic and linear
- Linear electron flow, the primary pathway, involves both photosystems and produces ATP and NADPH using light energy


- A photon hits a pigment and its energy is passed among pigment molecules until it excites P680
- An excited electron from P680 is transferred to the primary electron acceptor (we now call it P680⁺)




- P680⁺ is a very strong oxidizing agent
- H₂O is split by enzymes, and the electrons are transferred from the hydrogen atoms to P680⁺, thus reducing it to P680
- O₂ is released as a by-product of this reaction



- Each electron "falls" down an electron transport chain from the primary electron acceptor of PS II to PS I
- Energy released by the fall drives the creation of a proton gradient across the thylakoid membrane
- Diffusion of H⁺ (protons) across the membrane drives ATP synthesis




- In PS I (like PS II), transferred light energy excites P700, which loses an electron to an electron acceptor
- P700⁺ (P700 that is missing an electron) accepts an electron passed down from PS II via the electron transport chain




@ 2011 Peanon Education, Inc.

- Each electron "falls" down an electron transport chain from the primary electron acceptor of PS I to the protein ferredoxin (Fd)
- The electrons are then transferred to NADP⁺ and reduce it to NADPH
- The electrons of NADPH are available for the reactions of the Calvin cycle
- This process also removes an H⁺ from the stroma




@ 2011 Pearson Education, Inc.



Cyclic Electron Flow

- Cyclic electron flow uses only photosystem I and produces ATP, but not NADPH
- No oxygen is released
- Cyclic electron flow generates surplus ATP, satisfying the higher demand in the Calvin cycle




- Some organisms such as purple sulfur bacteria have PS I but not PS II
- Cyclic electron flow is thought to have evolved before linear electron flow
- Cyclic electron flow may protect cells from light-induced damage


A Comparison of Chemiosmosis in Chloroplasts and Mitochondria ATP SYMMUS to M Ariven by proto M gradienT

- Chloroplasts and mitochondria generate ATP by chemiosmosis, but use different sources of energy
- Mitochondria transfer chemical energy from food to ATP; chloroplasts transform light energy into the chemical energy of ATP
- Spatial organization of chemiosmosis differs between chloroplasts and mitochondria but also shows similarities

- In mitochondria, protons are pumped to the intermembrane space and drive ATP synthesis as they diffuse back into the mitochondrial matrix
- In chloroplasts, protons are pumped into the thylakoid space and drive ATP synthesis as they diffuse back into the stroma




- ATP and NADPH are produced on the side facing the stroma, where the Calvin cycle takes place
- In summary, light reactions generate ATP and increase the potential energy of electrons by moving them from H₂O to NADPH





Concept 10.3: The Calvin cycle uses the chemical energy of ATP and NADPH to reduce CO_2 to sugar

- The Calvin cycle, like the citric acid cycle, regenerates its starting material after molecules enter and leave the cycle
- The cycle builds sugar from smaller molecules by using ATP and the reducing power of electrons carried by NADPH

- Carbon enters the cycle as CO₂ and leaves as a sugar named glyceraldehyde 3-phospate (G3P)
- For net synthesis of 1 G3P, the cycle must take place three times, fixing 3 molecules of CO₂
- The Calvin cycle has three phases
 - Carbon fixation (catalyzed by rubisco)
 - Reduction
 - Regeneration of the CO₂ acceptor (RuBP)







Concept 10.4: Alternative mechanisms of carbon fixation have evolved in hot, arid climates

- Dehydration is a problem for plants, sometimes requiring trade-offs with other metabolic processes, especially photosynthesis
- On hot, dry days, plants close stomata, which conserves H₂O but also limits photosynthesis
- The closing of stomata reduces access to CO₂ and causes O₂ to build up
- These conditions favor an apparently wasteful process called photorespiration


Photorespiration: An Evolutionary Relic?


- In most plants (C₃ plants), initial fixation of CO₂, via rubisco, forms a three-carbon compound (3-phosphoglycerate)
- In photorespiration, rubisco adds O₂ instead of CO₂ in the Calvin cycle, producing a two-carbon compound
- Photorespiration consumes O₂ and organic fuel and releases CO₂ without producing ATP or sugar


- Photorespiration may be an evolutionary relic because rubisco first evolved at a time when the atmosphere had far less O₂ and more CO₂
- Photorespiration limits damaging products of light reactions that build up in the absence of the Calvin cycle
- In many plants, photorespiration is a problem because on a hot, dry day it can drain as much as 50% of the carbon fixed by the Calvin cycle

C₄ Plants

- **C**₄ **plants** minimize the cost of photorespiration by incorporating CO₂ into four-carbon phospho enolcompounds in mesophyll cells
- This step requires the enzyme **PEP** carboxylase
- PEP carboxylase has a higher affinity for CO₂ than rubisco does; it can fix CO₂ even when CO₂ concentrations are low
- These four-carbon compounds are exported to bundle-sheath cells, where they release CO₂ that is then used in the Calvin cycle








- In the last 150 years since the Industrial Revolution, CO₂ levels have risen greatly
- Increasing levels of CO₂ may affect C₃ and C₄ plants differently, perhaps changing the relative abundance of these species
- The effects of such changes are unpredictable and a cause for concern

CAM Plants

- Some plants, including succulents, use crassulacean acid metabolism (CAM) to fix carbon
- CAM plants open their stomata at night, incorporating CO₂ into organic acids
- Stomata close during the day, and CO₂ is released from organic acids and used in the Calvin cycle

Figure 10.21





Sugarcane



Pineapple 0 2011 Peanon Education, Inc.

The Importance of Photosynthesis: A Review

- The energy entering chloroplasts as sunlight gets stored as chemical energy in organic compounds
- Sugar made in the chloroplasts supplies chemical energy and carbon skeletons to synthesize the organic molecules of cells
- Plants store excess sugar as starch in structures such as roots, tubers, seeds, and fruits
- In addition to food production, photosynthesis produces the O₂ in our atmosphere

